

Package: BiTSLS (via r-universe)

May 24, 2026

Type Package

Title Bidirectional Two-Stage Least Squares Estimation

Version 0.1.0

Date 2025-05-26

Description Implements bidirectional two-stage least squares (Bi-TSLS) estimation for identifying bidirectional causal effects between two variables in the presence of unmeasured confounding. The method uses proxy variables (negative control exposure and outcome) along with at least one covariate to handle confounding.

License MIT + file LICENSE

Encoding UTF-8

LazyData true

Imports stats

RoxygenNote 7.3.1

Depends R (>= 3.5.0)

Repository <https://jiaqi-min.r-universe.dev>

Date/Publication 2025-05-28 08:00:05 UTC

RemoteUrl <https://github.com/jiaqi-min/bitls>

RemoteRef HEAD

RemoteSha b72bfac11c292320fc88e24f0c809aec34ebe873

Contents

Bi_TSLS	2
Index	3

Bi_TSLS

*Bidirectional Two-Stage Least Squares Estimation***Description**

Performs bidirectional two-stage least squares (Bi-TSLS) estimation to identify bidirectional causal effects between X and Y using proxy variables Z and W. Requires at least one covariate in addition to X, Y, Z, and W.

Usage

```
Bi_TSLS(data, R_w = 0, R_z = 0)
```

Arguments

data	A data frame containing at least these variables: <ul style="list-style-type: none"> • X: Treatment/exposure variable • Y: Outcome variable • Z: Negative control exposure (proxy variable) • W: Negative control outcome (proxy variable) • At least one additional covariate (required)
R_w	Sensitivity parameter for W (default = 0)
R_z	Sensitivity parameter for Z (default = 0)

Value

A list containing:

- beta_XY: Estimated effect of X on Y
- beta_YX: Estimated effect of Y on X

Examples

```
# Create example data with covariate
n <- 1000
data <- data.frame(
  X = rnorm(n),
  Y = rnorm(n),
  Z = rnorm(n),
  W = rnorm(n),
  V = rnorm(n) # At least one covariate is required
)

# Run BiTSLS
result <- Bi_TSLS(data)
print(result)
```

Index

Bi_TSLs, [2](#)